

SCHAUDER BASES UNDER UNIFORM RENORMINGS

A. J. GUIRAO AND P. HÁJEK

ABSTRACT. Let X be a separable superreflexive Banach space with a Schauder basis. We prove the existence of an equivalent uniformly smooth (resp. uniformly rotund) renorming under which the given basis is monotone.

1. INTRODUCTION

Let us start by recalling a classical notion of Schauder basis $\{e_n\}_{n=1}^\infty$ of a Banach space X , i.e. a sequence in X such that for every $x \in X$ there is a unique sequence of coefficients $\{a_n\}_{n=1}^\infty$, such that $x = \sum a_n e_n$. A Schauder basis gives rise to the canonical sequence of finite dimensional projections $P_n : X \rightarrow X$, $P_n(\sum_{i=1}^\infty a_i e_i) = \sum_{i=1}^n a_i e_i$. A well-known and useful result on Schauder bases claims that $\{\|P_n\|\}_n$ is a bounded sequence ([2]). The quantity $\text{bc}\{e_i\} = \sup_n \|P_n\|$ is called the basis constant. We say that the Schauder basis is monotone, if the value of its basis constant is 1. The boundedness result has an equivalent reformulation in the language of renormings. Namely, a separable Banach space with a Schauder basis can be equivalently renormed so that the basis becomes monotone. The proof of this equivalence is very easy, in fact the renorming is obtained via the formula $\|x\| = \sup_n \|P_n(x)\|$. Unfortunately, from the renorming point of view, $\|\cdot\|$ loses some subtler geometrical properties of the original norm. The question on the existence of "good" renormings, still making the given basis monotone, has received some attention in the past. For example, it is well-known that every separable Banach has an LUR renorming, in fact the collection of all equivalent LUR renormings is residual in the (metric) space of all equivalent renormings ([1]). It is therefore quite natural to expect that for every separable Banach space with a Schauder basis, there exists an equivalent LUR renorming making the basis monotone. This is indeed the case, and the proof follows along the lines of the original Kadec LUR renorming result (see [1]). Similar statements (folklore) are true also for Gateaux or uniformly Gateaux smooth case. On the

2000 *Mathematics Subject Classification.* 46B03.

Key words and phrases. Banach Spaces; Monotone Schauder Basis; Uniformly Fréchet Norms; Uniformly Rotund Norms.

First author supported by the grants MTM2005-08379 of MECD (Spain), 00690/PI/04 of Fundación Séneca (CARM, Spain) and AP2003-4453 of MECD (Spain), Second author supported by AV0Z10190503 and A100190502.

other hand, quite surprisingly, the situation with Fréchet smooth renormings is different. Recall that a separable Banach space has an equivalent Fréchet smooth renorming if and only if it has a separable dual. In this case, the set of Fréchet smooth renormings (whose dual norm is LUR) is again residual among all equivalent renormings, yet we have the following theorem.

Theorem 1. *Let X be a separable Banach space with a separable dual (in particular having an equivalent Fréchet smooth renorming), and a Schauder basis. Then X is reflexive iff for every Schauder basis of X there exists some Fréchet smooth renorming of X making the basis monotone.*

Proof. If X is reflexive, then X^* is separable and every Schauder basis of X is shrinking, so we can use the mentioned LUR result for the dual basis, in order to obtain the Fréchet smooth case for X (see [1] for details on the duality of renormings). On the other hand, by Proposition 8.34 of [2], if $\{e_i\}$ is a monotone basis of $(X, \|\cdot\|)$, and $\|\cdot\|$ is Fréchet smooth, then $\{e_i\}$ is a shrinking basis. Thus for spaces satisfying the second condition, every Schauder basis is shrinking. This condition is equivalent to reflexivity by a result of Zippin [5]. \square

As an immediate consequence, in every non-reflexive Banach space with a separable dual and a Schauder basis (such as c_0), there exists another Schauder basis which is not monotone under any Fréchet smooth renorming of X (in spite of the rich supply of equivalent Fréchet smooth renormings for such a space).

In the present note, we settle in the positive the case of uniformly Fréchet smooth (UF) and uniformly rotund (UR) renormings, answering a question of Godefroy (which appears also explicitly in [3]), communicated to us by Zizler. We would like to thank Václav Zizler for suggesting the problem to us, as well as for some useful remarks concerning this note.

1.1. Notation and basic definitions. We start by recalling some standard notions and definitions to be used throughout this note. The standard references for most of these notions are [2], or [1].

Definition 2. *Let B be a bounded, closed, convex and centrally symmetric subset of a linear space \mathbb{E} whose interior contains the origin. Then, we will say that B is a ball of \mathbb{E} .*

For every ball B of \mathbb{E} , there exists a norm in \mathbb{E} whose unit ball is precisely B . This norm is defined by the gauge functional, also known as Minkowski functional, and we will denote it, as it is usual in the literature, by $g(B, \cdot)$. We will also denote $S_B = \{x \in X : g(B, x) = 1\}$.

Regarding differentiability, we can define, for every ball B of \mathbb{E} , its modulus of smoothness.

Definition 3. Let B be a ball in a linear space \mathbb{E} . Then, its modulus of smoothness is the real function defined, for $t > 0$, by

$$\rho(B, t) := \sup \{ \xi(B, x, y, t) : g(B, x) = g(B, y) = 1 \}$$

where

$$\xi(B, x, y, t) = \frac{g(B, x + ty) + g(B, x - ty) - 2g(B, x)}{2}.$$

A norm $\| \cdot \|$ in Banach space is said to be *uniformly Fréchet differentiable*, shortly, UF, if its modulus of smoothness ρ satisfies the following condition:

$$\lim_{t \rightarrow 0} \frac{\rho(t)}{t} = 0.$$

Therefore we can say that, a ball B of a Banach space X is UF, if and only if the following condition holds:

$$\lim_{t \rightarrow 0} \frac{\rho(B, t)}{t} = 0.$$

Then, it is clear that B will be UF if and only if its associated norm by the gauge functional, $g(B, \cdot)$, is UF.

For every ball B , and $x \in \mathbb{E}$ satisfying $g(B, x) = 1$, let us denote by $H(B, x)$ the union of all the support hyperplanes to B at x . It is also useful the following modulus,

$$\bar{\rho}(B, t) := \sup \{ \xi(B, x, y, t) : g(B, x) = g(B, y) = 1 \text{ and } y + x \in H(B, x) \}.$$

It was shown by Figiel [4] that for each ball and for $t > 0$

$$\rho(B, t) \leq 16\bar{\rho}(B, t).$$

We will refer to this result as the Figiel lemma.

2. PRELIMINARY CONSTRUCTIONS

Let X be a Banach space endowed with a uniformly Fréchet differentiable norm $\| \cdot \|$, and a Schauder basis $\{e_i\}_{i=1}^{\infty}$. Our aim is to show that there exists an equivalent UF renorming of X which makes $\{e_i\}$ a monotone Schauder basis of X . Therefore, we will suppose from now on that the basis constant of the Schauder basis, $bc\{e_i\}$ is strictly greater than 1. Then, we can consider the norm $\|x\|_0 = \sup\{\|P_n(x)\| : n \in \mathbb{N}\}$, which is known to make the basis $\{e_i\}$ monotone (but of course not UF, in general).

Let us denote by B and B_0 the following balls:

$$B := \{x \in X : \|x\| \leq 1\},$$

$$B_0 := \{x \in X : \|x\|_0 \leq 1\}.$$

It is clear that

$$\| \cdot \| \leq \| \cdot \|_0 \leq bc\{e_i\} \| \cdot \|$$

and therefore

$$B_0 \subset B \subset bc\{e_i\}B_0.$$

Through this note $k \geq bc\{e_i\}$ is fixed. It is clear that the last two expressions are true replacing $bc\{e_i\}$ by k . For every natural number n , we will denote by \mathbb{E}_n the range of P_n , this is, $P_n X = \mathbb{E}_n$; by B'_n the image $P_n\left(\frac{1}{k}B\right)$; and for every $y \in \mathbb{E}_n$, by $B'_n(y)$ the set $y + B'_n$. In particular $B'_n(0) = B'_n$ is a ball of \mathbb{E}_n whose centre is the origin. For any $y \in \mathbb{E}_n$ the set $B'_n(y)$ is a translate of a ball of \mathbb{E}_n . We are going to define inductively a sequence $\{B_n\}_{n=1}^\infty$ of balls of \mathbb{E}_n as follows. Put $B_1 := B_0 \cap \mathbb{E}_1$. Having defined B_n for $n \geq 1$, we will define the set B_{n+1} as follows: First of all let us consider the set

$$\mathbb{B}_{n+1} := \{y \in \mathbb{E}_{n+1} : P_n(B'_{n+1}(y)) \subset B_n, B'_{n+1}(y) \cap B_0 \neq \emptyset\}$$

and then we finally set

$$B_{n+1} := \bigcup_{y \in \mathbb{B}_{n+1}} B'_{n+1}(y).$$

We can see, for example, the construction of B_2 in figure 1.




FIGURE 1

Lemma 4. *The family of sets $\{B_n\}_{n \in \mathbb{N}}$ is uniformly bounded in X , and for every n , the set B_n has non-empty interior with respect to the topology of \mathbb{E}_n .*

Proof. Let us fix $n \in \mathbb{N}$ and $x \in B_n$. Then there exist $y, z \in \mathbb{E}_n$ such that $x \in B'_n(y)$, $z \in B_0 \cap B'_n(y)$ and $P_{n-1}(B'_n(y)) \subset B_{n-1}$. It is clear that the vectors $x - y$ and $z - y$ lie in B'_n . Therefore, there exist $x', z' \in B$, such that $P_n(x') = k(x - y)$ and $P_n(z') = k(z - y)$, and then

$$\begin{aligned} \|x - y\| &= \frac{1}{k} \|P_n(x')\| \leq \frac{1}{k} \|P_n\| \|x'\| \leq 1, \\ \|z - y\| &= \frac{1}{k} \|P_n(z')\| \leq \frac{1}{k} \|P_n\| \|z'\| \leq 1. \end{aligned}$$

Since $z \in B_0$ and $B_0 \subset B$, we have $\|z\| \leq 1$. Therefore $\|x\| \leq 3$, this is $B_n \subset 3B \cap \mathbb{E}_n$. The second part of this proof is a direct application of the following lemma, since $B \cap \mathbb{E}_n \subset P_n(B)$. \square

Lemma 5. *For every natural number n , the set \mathbb{B}_n is non-empty. Moreover*

$$0 \in \bigcap_{n \geq 1} \mathbb{B}_n.$$

Proof. It is enough to show that $0 \in \mathbb{B}_n$ for any $n \in \mathbb{N}$. We will prove that $0 \in \mathbb{B}_n$ if and only if $B'_1 \subset B_1$, which will finish the proof, because it always holds

$$B'_1 = P_1 \left(\frac{1}{k} B \right) \subset P_1(B_0) \subset B_0 \cap \mathbb{E}_1 = B_1$$

since B_0 is a monotone ball. We will show the required equivalence by induction on n . For $n = 2$, 0 lies in \mathbb{B}_2 if and only if $P_1(B'_2) \subset B_1$, since B'_2 always intersects B_0 (the intersection contains at least the origin). But

$$P_1(B'_2) = P_1 \left(P_2 \left(\frac{1}{k} B \right) \right) = B'_1,$$

which finishes the proof of this case. If the statement is true for $n - 1 \geq 2$, then $0 \in \mathbb{B}_n$ if and only if $P_{n-1}(B'_n) \subset B_{n-1}$, since the intersection between B'_n and B_0 is always non-empty (it contains the origin). Besides

$$P_{n-1}(B'_n) = P_{n-1} \left(P_n \left(\frac{1}{k} B \right) \right) = P_{n-1} \left(\frac{1}{k} B \right) = B'_{n-1}.$$

Therefore $0 \in \mathbb{B}_n$ if and only if $0 \in \mathbb{B}_{n-1}$ which is equivalent to $B'_1 \subset B_1$. \square

The following lemma shows that the defined sets $\{B_n\}_n$ are balls in \mathbb{E}_n , i.e., their gauge functionals are norms in \mathbb{E}_n .

Lemma 6. *For every natural number n , the set B_n is a ball of \mathbb{E}_n .*

Proof. By lemma 4 it is clear that for every n , the set B_n is bounded and contains the origin as an interior point. For the remaining properties we will use induction.

It is clear that B_1 is closed, convex and centrally symmetric. Let us suppose that B_{n-1} do so for $n - 1 \geq 1$.

- a) We start proving that B_n is closed. Let us consider $x \in \overline{B_n}$. Then there exists a sequence $\{x_m\}_m$ in B_n converging to x . For every $m \in \mathbb{N}$ there exist y_m and z_m in \mathbb{E}_n such that $x_m \in B'_n(y_m)$, $z_m \in B_0 \cap B'_n(y_m)$ and $P_{n-1}(B'_n(y_m)) \subset B_{n-1}$. We can suppose that both $\{y_m\}$ and $\{z_m\}$ converge respectively to y and z in \mathbb{E}_n , since B_n is bounded. We want to show that $y \in B_n$.

Since $z_m \in B_0$ then $z \in B_0$, and it is clear that

$$\begin{aligned} \frac{1}{k} &\geq \lim_{m \rightarrow \infty} \|z_m - y_m\| = \|z - y\|, \\ \frac{1}{k} &\geq \lim_{m \rightarrow \infty} \|x_m - y_m\| = \|x - y\|. \end{aligned}$$

These equations mean, respectively, that $z \in B_0 \cap B'_n(y)$ and $x \in B'_n(y)$. Moreover, if we take $w \in B'_n(y)$, then $w_m := y_m + (w - y) \in B'_n(y_m)$ and $\{w_m\}_m$ converges to w . Therefore, since $P_{n-1}(w_m) \in B_{n-1}$ and by hypothesis B_{n-1} is closed, $P_{n-1}(w) \in B_{n-1}$, this is $x \in B_n$.

- b) Now we show that B_n is convex. Let us take x_1 and x_2 in B_n . For $i \in \{1, 2\}$ there exist $y_i \in \mathbb{B}_n$ such that $x_i \in B'_n(y_i)$, and $z_i \in B_0 \cap B'_n(y_i)$. Let us denote by $y = (y_1 + y_2)/2$, $z = (z_1 + z_2)/2$ and $x = (x_1 + x_2)/2$. Since $z_i - y_i \in B'_n$ for $i \in \{1, 2\}$ and B'_n is convex, then

$$z = \frac{z_1 + z_2}{2} = y + \frac{(z_1 - y_1) + (z_2 - y_2)}{2} \in B'_n(y).$$

Therefore $z \in B_0 \cap B'_n(y)$, since B_0 is also convex. Analogously we have $x \in B'_n(y)$. In order to prove that $x \in B_n$ we only need to see that $y \in \mathbb{B}_n$, so it is enough to show $P_{n-1}(B'_n(y)) \subset B_{n-1}$. In fact, if $w \in B'_n(y)$, then $w - y \in B'_n$ and for $i \in \{1, 2\}$ we have $y_i + (w - y) \in B'_n(y_i)$. Since $y_i \in \mathbb{B}_n$ for $i \in \{1, 2\}$, then $P_{n-1}(y_i + (w - y)) \in B_{n-1}$ for both $i = 1, 2$. Therefore

$$P_{n-1}(w) = \frac{1}{2} \left[P_{n-1}(y_1 + (w - y)) + P_{n-1}(y_2 + (w - y)) \right] \in B_{n-1},$$

since, by hypothesis, B_{n-1} is convex.

- c) Let $x \in B_n$. Then there exist $y \in \mathbb{B}_n$ such that $x \in B'_n(y)$, and $z \in B_0 \cap B'_n(y)$. It is clear that $z - y \in B'_n$ and, since B'_n is centrally symmetric, $y - z \in B'_n$. Therefore $-z = -y + (y - z) \in B'_n(-y)$. But B_0 is also centrally symmetric, and thus $-z \in B_0 \cap B'_n(-y)$. Analogously we can show that $-x \in B'_n(-y)$.

On the other hand, if we take $w \in B'_n(-y)$, then $w + y \in B'_n$, and, since B'_n is centrally symmetric, $-w - y \in B'_n$. Therefore $-w = y + (-w - y) \in B'_n(y)$. Since $P_{n-1}(B'_n(y)) \subset B_{n-1}$, and, by induction hypothesis, B_{n-1} is centrally symmetric, $P_{n-1}(w) \in B_{n-1}$. Hence, we have shown that $P_{n-1}(B'_n(-y)) \subset B_{n-1}$ and then $-y \in \mathbb{B}_n$. Therefore $-x \in B'_n(-y) \subset B_n$, this is, B_n is centrally symmetric. \square

In order to show that $\{B_n\}_n$ consists of UF-smooth balls, we need two auxiliary lemmas.

Lemma 7. *For any $n \in \mathbb{N}$, the modulus of smoothness of B'_n and B are related as follows:*

$$\rho(B'_n, \cdot) \leq \rho(B, \cdot).$$

Proof. Let us fix $n \in \mathbb{N}$ and $t > 0$. Let us also take x and y such that $g(B'_n, x) = g(B'_n, y) = 1$. In particular, since x and y lie on $S_{B'_n}$, there exist

$x', y' \in S_B$ such that $P_n(x') = kx$ and $P_n(y') = ky$. Let us note that

$$g\left(B'_n, P_n\left(\frac{x' \pm ty'}{kg(B, x' \pm ty')}\right)\right) \leq 1$$

This inequality is equivalent to $g(B'_n, x \pm ty) \leq g(B, x' \pm ty')$. Therefore

$$\xi(B'_n, x, y, t) \leq \xi(B, x', y', t) \leq \rho(B, t),$$

and taking suprema over $(x, y) \in S_{B'_n} \times S_{B'_n}$ we have the required result. \square

Lemma 8. *Let A and B be two balls in a linear space \mathbb{E} . Let $x \in \mathbb{E}$ such that $x + B \subset A$. Then, for any $z \in \mathbb{E}$ the following inequality holds:*

$$g(A, z) \leq g(B, z - x) + |1 - g(B, z - x)|g(A, x).$$

Proof. Let us note that the vector

$$\tilde{z} = x + \frac{z - x}{g(B, z - x)}$$

lies in the set A , see figure 2. Then, it is clear that

$$1 \geq g(A, \tilde{z}) = g\left(A, \frac{z - x(1 - g(B, z - x))}{g(B, z - x)}\right).$$




FIGURE 2

Therefore,

$$\begin{aligned} g(B, z - x) &\geq g(A, z - x(1 - g(B, z - x))) \\ &\geq g(A, z) - g(A, x(1 - g(B, z - x))). \end{aligned}$$

Thus we finally obtain

$$\begin{aligned} g(A, z) &\leq g(B, z - x) + g(A, x(1 - g(B, z - x))) \\ &= g(B, z - x) + |1 - g(B, z - x)|g(A, x). \end{aligned}$$

which concludes the proof. \square

Proposition 9. *For every $n \in \mathbb{N}$ the modulus of smoothness of B_n satisfies*

$$\rho(B_n, t) \leq 32\rho(B, 3kt).$$

Proof. Let us fix $n \in \mathbb{N}$, and $t > 0$. Let us also consider x and h points on S_{B_n} . By construction, there exists $y \in \mathbb{B}_n$ such that $g(B'_n, x - y) = 1$. Applying lemma 8 to B_n , B'_n and the point y , we can assure that

$$g(B_n, x \pm th) \leq g(B'_n, x \pm th - y) + [g(B'_n, x \pm th - y) - 1]g(B_n, y)$$

under the hypothesis $g(B'_n, x \pm th - y) \geq 1$. Therefore,

$$\begin{aligned} \xi(B_n, x, h, t) &\leq \xi(B'_n, x - y, h, t) + g(B_n, y)\xi(B'_n, x - y, h, t) \\ &= (1 + g(B_n, y))\xi(B'_n, x - y, h, t) \leq 2\xi(B'_n, x - y, h, t) \end{aligned}$$

If we denote by $\vartheta = g(B'_n, h)$, we have

$$\xi(B_n, x, h, t) \leq 2\xi(B'_n, x - y, h, t) = 2\xi(B'_n, x - y, \vartheta^{-1}h, \vartheta t) \leq 2\rho(B'_n, \vartheta t).$$

Since $\frac{1}{k}B \cap \mathbb{E}_n \subset B'_n$, by the monotonicity of the gauge functional we have

$$\begin{aligned} \vartheta &= g(B'_n, h) \leq g\left(\frac{1}{k}B \cap \mathbb{E}_n, h\right) = kg(B \cap \mathbb{E}_n, h) \\ &= kg(B, h) = k\|h\|. \end{aligned}$$

But, since $h \in B_n$, using lemma 4, we obtain $\|h\| \leq 3$.

Since the modulus of smoothness is a non-decreasing function, we finally have

$$\xi(B_n, x, h, t) \leq 2\rho(B'_n, \vartheta t) \leq 2\rho(B'_n, 3kt) \leq 2\rho(B, 3kt).$$

The last inequality follows directly from lemma 7. Taking suprema over x and h satisfying $g(B'_n, x \pm th - y) \geq 1$ we obtain that

$$\bar{\rho}(B_n, t) \leq 2\rho(B, 3kt).$$

The proof finishes applying Figiel's lemma. \square

3. UNIFORMLY FRÉCHET AND MONOTONE RENORMING

Now, we are ready to construct the new norm in X which will satisfy both uniformly Fréchet differentiability and monotonicity. In order to get it let us consider first some new sets. Let us fix $m \in \mathbb{N}$ and define

$$\tilde{B}_m := \bigcap_{n \geq m} (B_n \cap \mathbb{E}_m) = \left(\bigcap_{n \geq m} B_n \right) \cap \mathbb{E}_m.$$

We need to check if these new sets define, in fact, new norms in \mathbb{E}_m for each natural m , respectively.

Lemma 10. *The family of sets $\{\tilde{B}_m\}_{m \in \mathbb{N}}$ is uniformly bounded in X , and for every fixed m , the set \tilde{B}_m , with respect to the topology of \mathbb{E}_n , has non-empty interior. Moreover*

$$\frac{1}{k}B \cap \mathbb{E}_m \subset \tilde{B}_m \subset 3B.$$

Proof. Using the proof of lemma 4 it is clear that

$$\tilde{B}_m = \bigcap_{n \geq m} (B_n \cap \mathbb{E}_m) \subset \bigcap_{n \geq m} ((3B \cap \mathbb{E}_n) \cap \mathbb{E}_m) = 3B \cap \mathbb{E}_m \subset 3B.$$

On the other hand, by lemma 5 we know that

$$\tilde{B}_m = \bigcap_{n \geq m} (B_n \cap \mathbb{E}_m) \supset \bigcap_{n \geq m} \left(\left(\frac{1}{k}B \cap \mathbb{E}_n \right) \cap \mathbb{E}_m \right) = \frac{1}{k}B \cap \mathbb{E}_m.$$

□

Lemma 11. *For any natural number m , the set \tilde{B}_m is a ball in \mathbb{E}_m .*

Proof. By the previous lemma, these sets are bounded and have the origin as an interior point. It remains to show that these sets are closed, convex and centrally symmetric. However, these three properties are evident, because each \tilde{B}_m is the intersection of sets that are closed, convex and centrally symmetric. □

Proposition 12. *For any natural number m , the modulus of smoothness of \tilde{B}_m satisfies*

$$\rho(\tilde{B}_m, t) \leq 32\rho(B, 3kt).$$

Proof. For a fixed $m \in \mathbb{N}$, we consider $n \geq m$ and take $x \in B_{n+1} \cap \mathbb{E}_m$. Since $P_n(x) \in B_n$, we have that $x = P_m(x) = P_n(x) \in B_n$. This is, the sequence $\{B_n \cap \mathbb{E}_m\}_{n \geq m}$ is decreasing. Therefore it converges in the Hausdorff metric of \mathbb{E}_m to its intersection, this is, to \tilde{B}_m . It is then well-known, and standard to prove, that

$$(3.1) \quad \lim_{n \rightarrow \infty} g(B_n \cap \mathbb{E}_m, \cdot) = g(\tilde{B}_m, \cdot),$$

where the limit is understood as uniform convergence on bounded sets in \mathbb{E}_m .

Let us take x and y such that $g(\tilde{B}_m, x) = g(\tilde{B}_m, y) = 1$. We have

$$\begin{aligned} \xi(B_n, x, y, t) &= g(B_n, x) \xi \left(B_n, \frac{x}{g(B_n, x)}, \frac{y}{g(B_n, y)}, t \frac{g(B_n, y)}{g(B_n, x)} \right) \\ &\leq g(B_n, x) \rho \left(B_n, t \frac{g(B_n, y)}{g(B_n, x)} \right). \end{aligned}$$

Then by proposition 9

$$\xi(B_n, x, y, t) \leq g(B_n, x) 32\rho\left(B, 3kt \frac{g(B_n, y)}{g(B_n, x)}\right),$$

and taking limits, using equation (3.1) and the continuity of $\rho(B, \cdot)$, we have

$$\xi(\tilde{B}_m, x, y, t) \leq 32\rho(B, 3kt).$$

Taking suprema over x and y we obtain $\rho(\tilde{B}_m, t) \leq 32\rho(B, 3kt)$ as we wanted to show. \square

Finally, let us define the set

$$\tilde{B} := \bigcup_{m \in \mathbb{N}} \tilde{B}_m^{\|\cdot\|}.$$

Lemma 13. *For any natural number m , the following conditions hold:*

- (i) *For every $n > m$, $\tilde{B}_n \cap \mathbb{E}_m = \tilde{B}_m$.*
- (ii) *$\tilde{B}_n \subset \tilde{B}_m + \text{span}\{e_j : m < j \leq n\}$.*
- (iii) *$\tilde{B}_m = \tilde{B} \cap \mathbb{E}_m$.*

Proof. (i) Since the sequence $\{B_r \cap \mathbb{E}_m\}_{r \geq m}$ is decreasing, then

$$\tilde{B}_n \cap \mathbb{E}_m = \bigcap_{r \geq n} (B_r \cap \mathbb{E}_n) \cap \mathbb{E}_m = \bigcap_{r \geq n} (B_r \cap \mathbb{E}_m) = \bigcap_{r \geq m} (B_r \cap \mathbb{E}_m) = \tilde{B}_m.$$

- (ii) It is enough to show that $\tilde{B}_n \subset \tilde{B}_{n-1} + \text{span}\{e_n\}$. Let us take $x \in \tilde{B}_n$. Then $x \in B_r \cap \mathbb{E}_n$ for every $r \geq n$. Let us consider $r \geq n - 1$. Then $x \in B_{r+1} \cap \mathbb{E}_n$ and therefore x lies in $(B_r + \text{span}\{e_{r+1}\}) \cap \mathbb{E}_n$. This implies that $P_{n-1}(x) \in B_r \cap \mathbb{E}_{n-1}$. Since $r \geq n - 1$ is arbitrary, we have that $P_{n-1}(x)$ lies in \tilde{B}_{n-1} , this is, $x \in \tilde{B}_{n-1} + \text{span}\{e_n\}$.

- (iii) Clearly $\tilde{B}_m \subset \tilde{B} \cap \mathbb{E}_m$. On the other hand, let us take $x \in \tilde{B} \cap \mathbb{E}_m$. Then, there exists a sequence $\{x_n\}_n$ which converges to x and such that $x_n \in \tilde{B}_{m_n}$ for certain values $m_n \in \mathbb{N}$. Without loss of generality, we can reduce our proof to two cases:

- If $m_n \leq m$ for every $n \in \mathbb{N}$, then by the first item of this lemma, $x_n \in \tilde{B}_m$ and, since \tilde{B}_m is closed, $x \in \tilde{B}_m$.
- If $m_n > m$ for every $n \in \mathbb{N}$, by the second item of this lemma we have $P_m(x_n) \in \tilde{B}_m$. But, since $x \in \mathbb{E}_m$, we have that $\{P_m(x_n)\}_n$ converges to x , and therefore $x \in \tilde{B}_m$. \square

The following proposition summarizes the main properties of \tilde{B} . Condition (i) claims that it is an equivalent ball in X , (ii) implies that the basis $\{e_i\}$ is monotone with respect to the associated norm of \tilde{B} , and (iii) gives an estimate of the modulus of uniform smoothness.

Proposition 14. \tilde{B} is a ball of X , which satisfies the following conditions:

- (i) $\tilde{B} \subset 3B \subset 3k\tilde{B}$.
- (ii) For every $x \in \tilde{B}$ and for every natural number n , $P_n(x) \in \tilde{B}$.
- (iii) $\rho(\tilde{B}, t) \leq 32\rho(B, 3kt)$.

Proof. By lemma 13 (iii), the sequence $\{\tilde{B}_m\}_m$ is increasing. Therefore \tilde{B} is clearly convex and centrally symmetric. It follows from the definition of \tilde{B} that it is closed in X . In order to prove that \tilde{B} is a ball we just need to show that it contains the origin as an interior point, which is a consequence of item (i). Therefore we only need to show (i), (ii) and (iii).

- (i) By lemma 10 we have that $\tilde{B}_m \subset 3B$ for every $m \in \mathbb{N}$, therefore it is clear that $\tilde{B} \subset 3B$. On the other hand, let us take x such that $\|x\| < 1/k$. Since the sequence $\{P_m(x)\}_m$ converges to x , there exists m_0 , such that if $m \geq m_0$ then $\|P_m(x)\| \leq 1/k$. Since, by lemma 10, $B \cap \mathbb{E}_m \subset k\tilde{B}_m$, we have $P_m(x) \in \tilde{B}_m$ for every $m \geq m_0$, and then $P_m(x) \in \tilde{B}$ for every $m \geq m_0$. Therefore x lies in \tilde{B} . Then $sB \subset \tilde{B}$ for any $s < 1/k$. Taking closures $B \subset k\tilde{B}$.
- (ii) Let us take $x \in \tilde{B}$ and $m \in \mathbb{N}$. Let us also take $\{x_n\} \subset \tilde{B}$ in such a way $x_n \in \tilde{B}_{m_n}$. Without loss of generality, we can reduce our proof to two cases:
 - If $m_n \leq m$ for every $n \in \mathbb{N}$, then, by lemma 13 (i), $x_n \in \tilde{B}_m$, and then $P_m(x) = x$ lies in $\tilde{B}_m \subset \tilde{B}$.
 - If $m_n > m$ for every $n \in \mathbb{N}$, then, by lemma 13 (ii), $P_m(x_n) \in \tilde{B}_m$, thus $P_m(x)$ lies in $\tilde{B}_m \subset \tilde{B}$.
- (iii) From the previous results it is clear that the restriction of $g(\tilde{B}, \cdot)$ to \mathbb{E}_m coincides with $g(\tilde{B}_m, \cdot)$. Let us fix $t > 0$ and consider x and y such that $g(\tilde{B}, x) = g(\tilde{B}, y) = 1$. Let us denote $x_m = P_m(x)$, $y_m = P_m(y)$, $\theta_m = g(\tilde{B}_m, x_m)$ and $\vartheta_m = g(\tilde{B}_m, y_m)$. Then, by proposition 12,

$$\begin{aligned} \xi(\tilde{B}, x_m, y_m, t) &= \xi(\tilde{B}_m, x_m, y_m, t) = \theta_m \xi\left(\tilde{B}_m, \frac{x_m}{\theta_m}, \frac{y_m}{\vartheta_m}, t \frac{\vartheta_m}{\theta_m}\right) \\ &\leq \theta_m \rho\left(\tilde{B}_m, t \frac{\vartheta_m}{\theta_m}\right) \leq \theta_m 32\rho\left(B, 3kt \frac{\vartheta_m}{\theta_m}\right). \end{aligned}$$

Therefore

$$\begin{aligned} \xi(\tilde{B}, x, y, t) &= \lim_{m \rightarrow \infty} \xi(\tilde{B}, x_m, y_m, t) \leq \lim_{m \rightarrow \infty} \theta_m 32\rho\left(B, 3kt \frac{\vartheta_m}{\theta_m}\right) \\ &= 32\rho(B, 3kt). \end{aligned}$$

Taking suprema over x and y leads to the required conclusion. \square

As a simple corollary of this proposition we obtain the main results of our note.

Theorem 15. *Let X be a Banach space with a Schauder basis $\{e_i\}_{i=1}^{\infty}$ and a UF-norm $\|\cdot\|$. Then there exists an equivalent UF renorming $|||\cdot|||$ such that $\{e_i\}$ is a monotone Schauder basis of $(X, |||\cdot|||)$. Moreover, the moduli of uniform smoothness of $\|\cdot\|$ and $|||\cdot|||$ are equivalent (in the sense of [4]).*

Dualizing, we have the following result.

Theorem 16. *Let X be a separable superreflexive Banach space with a Schauder basis. Then there exists an equivalent UR renorming under which the basis is monotone.*

Proof. As X is superreflexive, by a well-known result of Enflo [6], there exist a UR-norm $\|\cdot\|$ on X . Since the basis is also shrinking, $(X^*, \|\cdot\|)$ is UF and $\{e_i^*\}$ is its Schauder basis. We can apply Theorem 15 to X^* , to obtain an equivalent UF-norm in X^* such that the basis $\{e_i^*\}$ is monotone. By duality, this means that $X^{**} = X$ has an equivalent UR-norm making the basis $\{e_i\}$ monotone. \square

REFERENCES

- [1] R. Deville, G. Godefroy, V. Zizler, Smoothness and renormings in Banach spaces, Pitman Monographs and Surveys **64**, Longman Ed (1993).
- [2] M. Fabian, P. Habala, P. Hájek, V. Montesinos, J. Pelant and V. Zizler, Functional analysis and infinite dimensional geometry, Canadian Math. Soc. Books, Springer Verlag, (2001).
- [3] M. Fabian, V. Montesinos and V. Zizler, Smoothness in Banach spaces. Selected problems, Rev. R. Acad. Cien. Serie A Mat. **100**, (2006), 101–125.
- [4] T. Figiel, On the moduli of convexity and smoothness, Studia Math. **56**, (1976), 121–155.
- [5] M. Zippin, A remark on bases and reflexivity in Banach spaces, Isr. J. Math. **6**, (1968), 74–79.
- [6] P. Enflo, Banach spaces which can be given an equivalent uniformly convex norm, Isr. J. Math **13**, (1972), 281–288.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, 30100 ESPINARDO (MURCIA), SPAIN

MATHEMATICAL INSTITUTE, AV ČR, ŽITNÁ 25, 115 67 PRAHA 1, CZECH REPUBLIC
E-mail address: `ajguirao@um.es` and `hajek@math.cas.cz`